



CONSENSUS STATEMENT

# Enhanced recovery after bariatric surgery: an Italian consensus statement

Giuseppe Marinari<sup>1</sup> · Mirto Foletto<sup>2</sup> · Carlo Nagliati<sup>3</sup> · Giuseppe Navarra<sup>4</sup> · Vincenzo Borrelli<sup>5</sup> · Vincenzo Bruni<sup>6</sup> · Giovanni Fantola<sup>7</sup> · Roberto Moroni<sup>8</sup> · Luigi Tritapepe<sup>9</sup> · Roberta Monzani<sup>10</sup> · Daniela Sanna<sup>11</sup> · Michele Carron<sup>12</sup> · Rita Cataldo<sup>13</sup>

Received: 17 August 2021 / Accepted: 31 December 2021  
© The Author(s) 2022

## Abstract

**Background** Enhanced recovery after bariatric surgery (ERABS) is an approach developed to improve outcomes in obese surgical patients. Unfortunately, it is not evenly implemented in Italy. The Italian Society for the Surgery of Obesity and Metabolic Diseases and the Italian Society of Anesthesia, Analgesia, Resuscitation and Intensive Care joined in drafting an official statement on ERABS.

**Methods** To assess the effectiveness and safety of ERABS and to develop evidence-based recommendations with regard to pre-, intra-, and post-operative care for obese patients undergoing ERABS, a 13-member expert task force of surgeons and anesthesiologists from Italian certified IFSO center of excellence in bariatric surgery was established and a review of English-language papers conducted. Oxford 2011 Levels of Evidence and U.S. Preventive Services Task Force Grade Definitions were used to grade the level of evidence and the strength of recommendations, respectively. The supporting evidence and recommendations were reviewed and discussed by the entire group at meetings to achieve a final consensus.

**Results** Compared to the conventional approach, ERABS reduces the length of hospital stay and does not heighten the risk of major post-operative complications, re-operations, and hospital re-admissions, nor does it increase the overall surgical costs. A total of 25 recommendations were proposed, covering pre-operative evaluation and care (7 items), intra-operative management (1 item, 11 sub-items), and post-operative care and discharge (6 items).

**Conclusions** ERABS is an effective and safe approach. The recommendations allow the proper management of obese patients undergoing ERABS for a better outcome.

**Keywords** Obesity · Bariatric Surgery · Anesthesia · Enhanced recovery after surgery

✉ Michele Carron  
michele.carron@unipd.it

<sup>1</sup> Bariatric Surgery Unit, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy

<sup>2</sup> Bariatric Surgery Unit, Azienda Ospedale Università Padova, Padua, Italy

<sup>3</sup> Department of Surgery, San Giovanni di Dio Hospital, Gorizia, Italy

<sup>4</sup> Department of Human Pathology, University of Messina, Messina, Italy

<sup>5</sup> Bariatric Surgery Unit, Humanitas Gavazzeni, Bergamo, Italy

<sup>6</sup> Bariatric Surgery Unit, Campus Bio Medico University of Rome, Rome, Italy

<sup>7</sup> Bariatric Surgery Unit, ARNAS, G. Brotzu Hospital, Cagliari, Italy

<sup>8</sup> Bariatric Surgery Unit, Policlinico Sassarese, Sassari, Italy

<sup>9</sup> Department of Anesthesia and Intensive Care, San Camillo-Forlanini Hospital, Sapienza University of Rome, Rome, Italy

<sup>10</sup> Department of Anesthesia and Intensive Care Units, Humanitas Research Hospital, Humanitas University Milan, Rozzano, Milan, Italy

<sup>11</sup> Emergency Department, Section of Anesthesiology and Intensive Care, ARNAS, G. Brotzu Hospital, Cagliari, Italy

<sup>12</sup> Department of Medicine-DIMED, Section of Anesthesiology and Intensive Care, University of Padua, Via V. Gallucci, 13, 35121 Padua, Italy

<sup>13</sup> Unit of Anesthesia, Intensive Care and Pain Management, Campus Bio Medico University of Rome, Rome, Italy

Bariatric surgery is the most effective treatment for morbid obesity. The demand for bariatric surgery continues to grow as the proportion of the population with obesity increases. Bariatric surgery is mainly optional in predominantly high-risk patients. Providing an effective, safe, and economically sustainable approach is challenging for physicians [1, 2].

Enhanced Recovery after Surgery (ERAS) is a modern approach to reduce perioperative stress and help patients recover more quickly following surgery, leading to improved outcomes in postoperative morbidity and length of hospital stay (LOS). ERAS, implemented initially for elective colorectal surgery, has been extended over time to many other surgeries [3], including bariatric surgery [4]. However, the ERAS after bariatric surgery (ERABS) guidelines published in 2016 [4] relied heavily on the evidence and recommendations developed in no-bariatric surgery settings. Afterward, an increasing number of peer-reviewed papers about ERABS have been published, showing the favorable outcomes observed in ERAS.

Despite the evidence, some resistance to adopting ERABS protocols remains in bariatric surgery centers in Italy. The Italian Ministry of Health's 2018 Annual Report on hospital admission and discharge events reported a total of 25,424 surgical procedures for obesity, with an average LOS of 3.9 days [5]. So, if, on the one hand, this data suggests that the conventional approach is largely adopted in bariatric surgery, on the other, it leaves considerable opportunity for improving bariatric surgery outcomes by adopting the ERABS approach.

## Materials and methods

The Italian Society of Surgery of Obesity and Metabolic Diseases (SICOB) and the Italian Society of Anesthesia, Analgesia, Resuscitation and Intensive Care (SIAARTI) undertook a process for drafting an official joint statement on ERABS. The two societies appointed a 13-member expert task force, which met first in October 2019 to define the scope and methods of the project. It was decided that the primary objective was to summarize available evidence supporting ERABS, focusing on the efficacy and safety outcomes. The secondary objective was to review the importance of the individual ERABS protocol items.

Four main areas for investigation were identified: outcomes for the primary endpoint and preoperative, intraoperative, and postoperative care for the secondary endpoint. Corresponding subcommittees were appointed to systematically review ERABS topics, grade the levels of evidence, and propose specific recommendations with supporting evidence for each topic.

A systematic PubMed search of English-language papers published from 1999 to April 2020 was performed

combining the following terms: “obesity” and “surgery” or “bariatric surgery” or “fast track” or “enhanced recovery” or “perioperative care” or “perioperative management” or “sleeve gastrectomy” or “gastric bypass.” The levels of evidence were assessed using Oxford 2011 Levels of Evidence (Table 1) [6]. To rate the strength of recommendations, grade definitions based on the U.S. Preventive Services Task Force were used (Tables 2 and 3) [7].

Key issues were discussed at a meeting in January 2020, after which a comprehensive document was circulated and then revised. The panel concluded its work with a Consensus Meeting in July 2020, where supporting evidence and recommendations were reviewed and discussed by the entire group to achieve a final consensus. Subsequently, a draft report was prepared and sent to the panel for comments and modification. Each author approved the final version before it was submitted. Ethics approval and consent were not required for this type of study.

## Results

The evidence and recommendations concerning the primary safety and efficiency endpoints are summarized in Table 4. A total of 25 recommendations were proposed, covering pre-operative evaluation and care (7 items), intra-operative management (1 item, 11 sub-items), and post-operative care and discharge (6 items). The levels of evidence and recommendations for each item in the ERABS protocol are summarized in Table 4.

### Effectiveness and safety of ERABS

Compared to the conventional approach, ERABS reduces the LOS [8–10], independently of the type of surgical procedure performed [11, 12]. The more successful ERABS items the multidisciplinary team adopts, the greater the likelihood of a reduced post-operative LOS [13].

Also, compared to the standard approach, ERABS does not have an increased risk for major postoperative complications, reoperations, and hospital readmissions [8–12, 14]. However, it also does not reduce the number of major complications [8–12, 14]. ERABS, moreover, seems to reduce the total surgical costs compared to the standard approach [8, 15].

### Preoperative counseling

Preoperative information and counseling are key items for managing the expectations of patients and preparing them for early discharge [16]. Counseling was one of the most frequently used items in a retrospective multicenter study [17] and one of the key items in 11 of the 13 studies considered in

**Table 1** Grading of level of evidence (from Oxford Centre for Evidence-Based Medicine 2011 Level of Evidence) [6]

Question	Step 1 (Level 1*)	Step 2 (Level 2*)	Step 3 (Level 3*)	Step 4 (Level 4*)	Step 5 (Level 5*)
How common is the problem?	Local and current random sample surveys (or censuses)	Systematic review of surveys that allow matching to local circumstances**	Local non-random sample**	Case-series**	n/a
Is this diagnostic or monitoring test accurate? (Diagnosis)	Systematic review of cross sectional studies with consistently applied reference standard and blinding	Individual cross sectional studies with consistently applied reference standard and blinding	Non-consecutive studies, or studies without consistently applied reference standards**	Case-control studies, or 'poor' or non-independent reference standard**	Mechanism-based reasoning
What will happen if we do not add a therapy? (Prognosis)	Systematic review of inception cohort studies	Inception cohort studies	Cohort study or control arm of randomized trial*	Case-series or case-control studies, or poor quality prognostic cohort study**	n/a
Does this intervention help? (Treatment Benefits)	Systematic review of randomized trials or n-of-1 trials	Randomized trial or observational study with dramatic effect	Non-randomized controlled cohort/follow-up study**	Case-series, case-control studies, or historically controlled studies**	Mechanism-based reasoning
What are the COMMON harms? (Treatment Harms)	Systematic review of randomized trials, systematic review of nested case-control studies, n of-1 trial with the patient you are raising the question about, or observational study with dramatic effect	Individual randomized trial or (exceptionally) observational study with dramatic effect	Non-randomized controlled cohort/follow-up study (post-marketing surveillance) provided there are sufficient numbers to rule out a common harm. (For long-term harms the duration of follow-up must be sufficient.)**	Case-series, case-control, or historically controlled studies**	Mechanism-based reasoning
What are the RARE harms? (Treatment Harms)	Systematic review of randomized trials or n-of-1 trial	Randomized trial or (exceptionally) observational study with dramatic effect			
Is this (early detection) test worthwhile? (Screening)	Systematic review of randomized trials	Randomized trial	Non-randomized controlled cohort/follow-up study**	Case-series, case-control, or historically controlled studies**	Mechanism-based reasoning

\*Level may be graded down on the basis of study quality, imprecision, indirectness (study PICO does not match questions PICO), because of inconsistency between studies, or because the absolute effect size is very small; Level may be graded up if there is a large or very large effect size

\*\*As always, a systematic review is generally better than an individual study

**Table 2** Grading of quality of evidence (from US Preventive Services Task Force) [7]

Grade	Description
A	The available evidence usually includes consistent results from multitude of well-designed, well-conducted, studies in representative care populations. These studies assess the effects of the service on the desired health outcomes. Because of the precision of findings, this conclusion is, therefore, unlikely to be strongly affected by the results of future studies. These recommendations are often based on direct evidence from clinical trials of screening, treatment, or behavioral interventions. High-quality trials designed as “pragmatic” or “effectiveness” trials are often of greater value in understanding external validity
B	The available evidence is sufficient to determine the effects of the service on targeted health outcomes, but confidence in the estimate is constrained by factors such as the number, size, or quality of individual studies in the evidence pool; some heterogeneity of outcome findings or intervention models across the body of studies; mild-to-moderate limitations in the generalizability of findings to routine care practice. As more information becomes available, the magnitude or direction of the observed effect could change, and this change may be large enough to alter the conclusion
C	The available evidence is insufficient to assess effects on health outcomes. Evidence is insufficient because of the very limited number or size of studies Inconsistency of direction or magnitude of findings across the body of evidence; critical gaps in the chain of evidence; findings are not generalizable to routine care practice; a lack of information on prespecified health outcomes; lack of coherence across the linkages in the chain of evidence. More information may allow an estimation of effects on health outcomes

**Table 3** Grading of the strength of recommendation (from U.S. Preventive Services Task Force) [7]

Grade	Definition	Suggestion for practice
A	The USPSTF recommends the service. There is high certainty that the net benefit is substantial	Offer or provide this service
B	The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial	Offer or provide this service
C	The USPSTF recommends selectively offering or providing this service to individual patients based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small	Offer or provide this service for selected patients depending on individual circumstances
D	The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits	Discourage the use of this service
I	The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined	Read the clinical considerations section of USPSTF Recommendation Statement. If the service is offered, patients should understand the uncertainty about the balance of benefits and harms

a systematic review [8]. A retrospective study on the safety of an early postoperative discharge following bariatric surgery identified the receipt of preoperative information on early ambulation and refeeding, as well as pain and postoperative nausea and vomiting (PONV) management, as one of the more important protocol items of the ERABS pathway [18].

### Preoperative patient optimization

Smoking cessation for at least four weeks reduced postoperative surgical and pulmonary complications by 41% [19], supporting the argument that smoking should be discontinued for at least four weeks before bariatric surgery [20]. Physicians should inform patients of the increased risk of morbidity and mortality in smokers. This risk decreases the longer smoking is ceased before surgery [21].

Preoperative weight loss reduces liver volume and may technically facilitate the operation [22]. However, whether weight loss reduces postoperative complications remains controversial [21, 23]. A Swedish registry study reported a decrease in complications after gastric bypass surgery [24].

Optimizing the preoperative fasting blood glucose level through diet, physical activity, and pharmacotherapy is mandatory [20, 21, 23]. A value greater than 180 mg/dl was associated with increased perioperative complications and mortality [20].

Obstructive sleep apnea (OSA) carries an increased risk of postoperative cardiorespiratory complications [25]. Although this aspect has been debated in bariatric surgery [26, 27], increased complications and LOS have been observed in OSA patients who underwent bariatric surgery [28–30]. The use of noninvasive ventilation (NIV) (e.g., continuous positive airway pressure [CPAP]) was reported to

**Table 4** Effectiveness, safety, and items of Enhanced Recovery after Bariatric Surgery (ERABS) compared to standard approach

	Evidence		Strength of Recommendation	Expert task force statement
	Level	Quality		
<i>Effectiveness and safety of ERABS</i>				
Length of hospital stay	1	A	A	ERABS reduces the duration of hospital stays
Risk of complications	1	A	A	ERABS is a safe approach for obese patients
Cost of surgery	2	B	A	Adopting an ERABS protocol does not increase the cost of surgery
<i>Items ERABS. Preoperative care</i>				
Information and counseling	2	B	A	The information provided to the patient should not be limited to what is required for informed consent for both surgery and anesthesia; it should be adequate to provide realistic expectations of the ERABS approach
Patient optimization	1	A	A	Pre-operative optimization through smoking cessation, weight loss, blood glucose control, and the use of non-invasive ventilation (when indicated) is recommended in ERABS
Fasting	1	A	A	Clear liquids and solid food are recommended up to 2 h and 6 h, respectively, prior to the induction of anesthesia in ERABS
PONV prophylaxis	1	A	A	Strategies aimed at minimizing the risk of post-operative nausea and vomiting after general anesthesia are recommended for better patient outcomes in ERABS
Venous thromboembolism prophylaxis	2	B	A	Multimodal venous thromboembolism prophylaxis, including early patient mobilization, is recommended in ERABS
Antibiotic prophylaxis	2	B	A	Pre-operative intravenous antibiotic prophylaxis is recommended in ERABS
Monitoring	1	A	A	Proper perioperative monitoring is recommended in ERABS
<i>Items ERABS. Intraoperative care</i>				
Standardized anesthesia protocol	1	A	A	A standardized anesthesia approach is recommended in order to optimize outcomes in ERABS
1-Airways management	1	A	A	A careful airways assessment is recommended in ERABS
2-Preoxygenation	1	A	A	An adequate preoxygenation performed in ramped position is recommended in ERABS
3-Tracheal intubation	1	A	A	Proper airway management in order to minimize difficulties is recommended in ERABS
4-General anesthesia	1	A	A	General anesthesia is the anesthesiologic approach of choice in ERABS
5-Neuromuscular blockade	1	A	A	Proper neuromuscular blockade management is recommended in ERABS
6-Analgesia opioid sparing	1	A	A	Opioid-sparing or opioid-free anesthesia is recommended in ERABS
7-Multimodal analgesia	1	A	A	Multimodal analgesia is recommended in ERABS to optimize pain control after surgery and to reduce or eliminate the use of opioids in the post-operative period
8-Locoregional anesthesia	1	A	A	Locoregional anesthesia supports and complements general anesthesia in ERABS
9-Protective lung ventilation	1	A	A	Protective mechanical lung ventilation during general anesthesia is recommended in ERABS
10-Goal-directed fluid therapy	3	B	A	Proper perioperative fluid management is recommended. Goal-directed fluid therapy should be considered in ERABS
11-Protected extubation	1	A	A	Extubation should be performed on an awake patient in the ramped position in ERABS
<i>Items ERABS. Postoperative care</i>				
Nasogastric tube	1	A	A	Routine placement of the SNG does not improve outcomes in ERABS
Abdominal drainage	2	B	A	Routine use of abdominal drainage should be discontinued in ERABS
Bladder catheter	4	C	A	Routine use of bladder catheters should be abandoned in ERABS

**Table 4** (continued)

	Evidence		Strength of Recommendation	Expert task force statement
	Level	Quality		
Early mobilization	3	B	A	Early post-operative mobilization is recommended in ERABS
Early re-feeding	1	A	A	Early post-operative resumption of oral feeding is recommended in ERABS
Early discharge	1	A	A	Early discharge of the patient is recommended in ERABS. Adoption and verification of a discharge checklist upon discharge are recommended in ERABS

improve the preoperative cardiometabolic profile [31] and the postoperative respiratory function [21, 23, 32]. It may be safely adopted for patients undergoing bariatric surgery [32]. Even if postoperative care in a monitored setting may be considered for high-risk obese patients undergoing surgery [20, 33], patients with severe OSA and/or home CPAP selected for ERABS do not require a routine planned postoperative admission into the intensive care unit [20, 21, 27].

### Preoperative fasting

Properly managing preoperative fasting may be important to minimize perioperative stress. Maintaining homeostasis avoids or reduces catabolism and related proteolysis, asthenia, or cellular dysfunction [34–37]. Clear liquids can be taken up to 2 h before surgery and solid food up to 6 h before the induction of anesthesia [3, 38–41]. No strong evidence supports preoperative oral carbohydrate loading in bariatric surgery. Furthermore, there is some reluctance to adopt preoperative oral carbohydrate loading in patients suffering from diabetes or metabolic syndrome [42]. Absolute or prolonged preoperative fasting is no guarantee of a secretion-free stomach at the induction of anesthesia [42].

### Postoperative nausea and vomiting prophylaxis

General anesthesia is associated with an increased risk of PONV [43]. The prevention of PONV in ERABS is recommended [4], and it is consistent with the guidelines for the general surgical population [43]. The polypharmacological approach to PONV prophylaxis is preferable to monotherapy [20, 43] and was reported to decrease the incidence of PONV and the postoperative use of antiemetics, opioid analgesics, and liquid infusion [44].

### Venous thromboembolism prophylaxis

The incidence of deep vein thrombosis and pulmonary embolism after bariatric surgery is approximately 0.1–0.5% and, in the vast majority of cases, occurs after the patient is discharged [21, 45, 46]. A multimodal venous

thromboembolism prophylaxis via chemoprophylaxis, mechanical aids, and/or patient mobilization is suggested [4, 20, 21, 47–49]. It is reported as a key item in ERABS [12, 34, 39, 50–61]. Chemoprophylaxis with low-molecular-weight heparin (LMWH) after surgery and home discharge is considered effective [20, 48, 52, 53] and is associated with a lower bleeding risk [52]. However, there is insufficient evidence to recommend a specific dose and duration of the LMWH treatment [20, 47–49, 52]. Early ambulation in the postoperative period is considered a useful component of multimodal venous thromboembolism prophylaxis [20, 47, 49].

### Antibiotic prophylaxis

The incidence of surgical site infections in obese patients varies from 1 to 21.7%, depending on the procedure type [62]. The use of preoperative antibiotic prophylaxis is then recommended [20, 21, 23]. It should follow the standard guidelines for perioperative antibiotic prophylaxis [63]. The most frequently used antibiotic is cefazolin (dosage 1–4 g), with clindamycin recommended as an alternative in allergic patients [54, 64]. A dose adjustment based on the patient's weight compared to a fixed dose of 2 g administered intravenously before surgical incision is preferable [65, 66]. Higher dosages (cefazolin 3 g) in patients weighing > 120 kg should be considered [20]. Literature does not support prophylactic vancomycin or cefoxitin [64]. Intestinal preparation by antibiotic prophylaxis (whether or not combined with mechanical preparation) is not recommended in bariatric surgery [60]. Preoperative antibiotic prophylaxis, administered intravenously at the induction of anesthesia or 30–60 min before surgical incision, is reported to be one of the important items in ERABS [12, 17, 60, 67].

### Monitoring

The standard for anesthesia monitoring should be ensured in the perioperative period [4, 20, 33, 68]. In obese patients, anesthesia depth monitoring ensures a more accurate induction of anesthesia with propofol and reduces the risk of



awareness from inadequate dosing [69], as well as during intravenous and inhalational anesthesia [70]. The monitoring of the neuromuscular function helps reduce the risk of postoperative respiratory complications [71]. Temperature monitoring reduces the risk of hypothermia, improving postoperative recovery [72].

### Standardized anesthesia approach

The standardized ERABS approach showed favorable outcomes compared to a non-standardized one [8–14]. Advances in the perioperative care of obese patients translated into a standardized anesthesiological approach have been proven to be effective and safe [73] in the ERABS context as well [39].

### Airway management

In obese surgical patients, airway management can present challenges. Difficult mask ventilation has been reported in 8.8% of obese patients, and 11% of those have morbid obesity. Difficult intubation has been reported in 3.3–16.7% of patients [20]. Difficult airway management predictors include males, OSA, increased waist-to-hip ratio ( $> 0.8$  in women and  $> 0.9$  in men), BMI ( $> 50 \text{ kg/m}^2$ ), and neck circumference ( $> 41 \text{ cm}$  in women and  $> 43 \text{ cm}$  in men) [20, 33, 74–76]. OSA risk stratification is then suggested using validated questionnaires, such as STOP-BANG, and reserving the time-consuming and expensive polysomnography for patients at high risk of severe OSA (e.g., STOP-BANG  $\geq 5$ ) [3, 21, 23, 28–31, 33, 77–81]. The ramp position improves the likelihood of successful airway management in obese patients [20, 23].

### Pre-oxygenation

Adequate pre-oxygenation aiming for end-tidal oxygen concentrations of  $\geq 90\%$  before the induction of general anesthesia is suggested. Pre-oxygenation has been reported as important in prolonging safe apnea times following general anesthesia induction [82, 83]. The NO DESAT (nasal oxygen during efforts securing a tube) technique, which uses a simple nasal cannula with standard cold dry oxygen may be considered [82]. Pre-oxygenation using a high-flow nasal cannula or positive pressure by CPAP/NIV seems more effective than the standard approach [82, 84–91] and may be beneficial for high-risk obese patients [20, 32, 33, 82].

### Tracheal intubation

An appropriate planned approach is recommended for airway management in obese patients [3, 20, 33]. In obese patients, videolaryngoscopes compared with Macintosh

laryngoscopes increases the likelihood of successful intubation on the first attempt at laryngoscopy [92]. This result seems to be ensured more by the use of a videolaryngoscope with a tracheal tube guide than one without [93].

Second-generation extraglottic devices were recommended as rescue devices for pulmonary oxygenation/ventilation in the case of difficult airway management and possible fibroscope-guided intubation [94–98]. The fibroscopic/endoscopic technique was essential when intubating conscious patients [99]. In awake patient, video laryngoscopy has been suggested as a valid alternative to fiberoptic intubation in experienced practitioners [100–102].

### General anesthesia

General anesthesia is the approach of choice in ERABS [3, 4, 20, 33]. There is no evidence supporting the superiority of inhalation versus intravenous anesthesia [3, 4, 20, 33]. Anesthesia strategies based on short-acting, low-accumulation drugs that promote rapid recovery from general anesthesia are suggested [4, 20, 33]. Desflurane has been associated with faster postoperative awakening and recovery than other inhalational anesthetics and propofol [103–106]. Intravenous anesthesia with propofol has demonstrated a lower risk of PONV in the general population [107] and in obese surgical patients [108]. PONV prophylaxis has been observed to reduce the risk of PONV incidence [109], particularly with inhalational anesthesia [20].

### Analgesia and opioid

In the general surgical population, opioid use was associated with an increased risk of PONV [110], as well as upper airway obstruction and hypoventilation [111]. In obese patients, intraoperative opioid use was associated with an increased risk of PONV [112] and postoperative respiratory complications [113]. *Opioid-sparing* or *opioid-free* anesthesia should be preferred when managing obese surgical patients [20, 33] under ERABS [4], as it is associated with a lower incidence of PONV [112].

Opioids with rapid elimination kinetics, such as remifentanyl, have demonstrated faster postoperative awakening and recovery of respiratory functions in general surgical populations compared to other opioids [114]. In obese patients, remifentanyl has demonstrated reduced recovery time from general anesthesia, respiratory complications, and LOS [115]. Patient-controlled analgesia has also been successfully used in obese surgical patients [116] and should be preferred over continuous infusion in the postoperative period [3, 20].

## Multimodal analgesia

Multimodal analgesia as a pain control strategy using analgesics (e.g., nonsteroidal anti-inflammatory drugs, paracetamol) in combination with analgesic adjuvants and local anesthetics is recommended in ERABS to optimize pain control after surgery and to reduce or eliminate the postoperative use of opioids [4, 20, 33, 34, 117] and related adverse events [118]. Adjuvant analgesics, such as ketamine [119, 120], clonidine [119, 121], dexmedetomidine [121–123], magnesium [123–125], lidocaine [126], pregabalin [127, 128], and gabapentin [129], alone or in combination, significantly reduced postoperative pain compared with controls.

## Locoregional analgesia

Intraperitoneal local anesthetics [130–133] and the ultrasound-guided TAP (transversus abdominis plane) block [134–136] showed a significant effect on reducing pain scores at recovery from general anesthesia [137, 138]. The benefit may be increased by the infiltration of trocar insertions and wound closures with local anesthetics at the end of the surgical procedure [20, 135]. Notably, these pain control benefits did not affect the LOS [130–136]. Epidural analgesia may be considered in selected patients [139].

## Neuromuscular blockade

Neuromuscular blockade (NMB) is suggested at the induction of general anesthesia to facilitate airway management and subsequent pulmonary ventilation [20, 33, 82]. Deep compared to moderate NMB optimizes the surgical field view, reduces procedural complications in laparoscopic surgery [140], and is associated with less postoperative pain [140, 141]. Sugammadex compared to cholinesterase inhibitors provides a more rapid and predictable recovery of rocuronium-induced NMB [142, 143] and is associated with less pain and PONV in the postoperative period [141, 144] and faster discharge to the surgical ward [143, 144]. Quantitative monitoring and complete recovery of the neuromuscular function at the end of surgery are highly recommended [4, 20, 33].

## Protective ventilation

Protective mechanical lung ventilation should be preferred for obese patients undergoing general anesthesia for bariatric surgery [20] because it is associated with reduced postoperative respiratory complications, LOS, and mortality in both non-obese and obese surgical patients [145–147].

Slow abdominal insufflation with a maximum intra-abdominal pressure of less than 15 mmHg is advised, when possible, during laparoscopy [20] to favor mechanical lung

ventilation [20, 148]. This strategy, combined with limiting surgical time, is associated with a reduction in the risk of postoperative respiratory complications [148–150].

## Goal-directed fluid therapy

Proper perioperative fluid management avoiding overhydration helps to minimize the risk of PONV, postoperative complications, and prolonged LOS [151–153]. A goal-directed fluid therapy (GDFT) has been suggested as an adequate strategy to reduce these risks [154], even in bariatric surgery [155–157]. In bariatric surgery, excessive fluid should be avoided, and GDFT should be considered a useful strategy [155–157]. A GDFT guided by noninvasive indices, such as the Pleth Variability Index, may be a more acceptable monitoring option to GDFT based on invasive methods [155] and, consequently, more widely used in bariatric surgery, even in the ERABS context [155–157]. Postoperative fluid infusions should be discontinued as soon as possible, with preference given to the enteral route [60, 153, 158]. Intraoperative hypotension (mean arterial pressure of  $\leq 65$  mmHg), even for a few minutes, is a predictor of renal and myocardial damage [159]. It should thus be avoided or promptly treated in the perioperative period [154, 159].

## Protected extubation

The extubation should be performed on an awake patient in the ramped position and/or reverse Trendelenburg position, which improves lung volume, oxygenation, and respiratory mechanics [20, 33, 160, 161]. Oxygen therapy with nasal goggles or HFNC was associated with a reduced risk of post-extubation desaturation [160] and reintubation [162]. CPAP or NIV was recommended for awakening moderate-severe OSA patients or those suffering from the obese hypoventilation syndrome who are already receiving home treatment or who will require opioid therapy postoperatively [20, 32, 163]. CPAP and NIV do not appear to negatively affect the outcome of the surgical procedure [32].

## Nasogastric tube

In abdominal surgery, avoiding the routine use of nasogastric tubes (NGTs) results in a faster recovery of the bowel function, a decrease in pulmonary complications, and a shorter LOS without any associated increase in anastomotic dehiscence [164]. In ERABS, compared to the conventional approach, avoiding NGT use demonstrated better postoperative recovery without an increase in complications [11–13, 41], reducing postoperative pain and PONV, promoting early mobilization and resumption of liquid diet, and resulting in better compliance at discharge [38, 39, 55]. Society positions



in favor of abandoning, when possible, the routine use of postoperative NGT are already available [3, 54].

### Abdominal drainage

Many gastrointestinal surgeries can be performed safely without prophylactic drainage [165]. In bariatric surgery, the evidence is limited [166]. A retrospective study of a gastric bypass population reported no difference in anastomotic dehiscence and reintervention rates in patients receiving abdominal drainage compared to those who are not [167]. Avoiding abdominal drainage was found to be a key item in the ERABS [3, 11, 12, 14, 17, 19, 59, 67, 168]. Society positions in favor of abandoning the routine use of abdominal drainage, when possible, are available [3, 54].

### Bladder catheter

Avoidance or early removal of bladder catheters resulted in early mobilization and prevented urinary tract infection [169–171]. In ERABS, compared to the conventional approach, avoiding bladder catheters was associated with early mobilization and reduced LOS, readmission rate, and minor complications, such as urinary tract infection [3, 10, 14, 18, 50, 67, 168, 172]. Society positions in favor of abandoning the routine use of bladder catheter, when possible, are available [54].

### Early mobilization

Early postoperative mobilization is recommended in obese patients undergoing bariatric surgery [4, 11, 14, 20, 21, 33]. Independent mobilization for at least 4 h during the first 24 h following surgery was significantly associated with a lower rate of postoperative complications and a shorter LOS [173]. Early postoperative mobilization was reported as a key element of better postoperative recovery in ERABS [17–20, 29, 39, 54–59, 67, 168, 172], and adequate independent mobilization was a condition for a home discharge [14, 18, 21, 39, 54, 56, 58, 59, 67].

### Early refeeding

The advantage of early oral refeeding has been reported in both ERAS [3] and ERABS [39–41, 173] and seems particularly associated with an earlier restoration of bowel function, faster wound healing, less infection, and lower risk of postoperative complications [36, 37, 174]. Early oral refeeding was associated with a reduced LOS and mortality [44, 60]. Conversely, prolonged postoperative fasting was associated with thirst, emotional fixation on food, and a phobia about the reintroduction of food [36, 37, 174].

### Discharge

Discharge on postoperative day 1 or 2 was not associated with an increase in the complication rate, readmission, or telephone consultations in both ERAS [3, 18] and ERABS [11, 12, 175], particularly in the absence of significant comorbidities [176, 177]. Caution should be taken in the presence of suspected clinical conditions (e.g., tachycardia) and/or abnormal level of serological markers (e.g., C-reactive protein, procalcitonin) as predictors of risk of postoperative complications [178–181]. However, there is no consensus on which serological markers should be assessed and at which postoperative time point [182–185]. Routine postoperative contrast imaging examination increased costs and LOS [186, 187] and was not considered to be a reliable assessment of postoperative complications [188–190].

Discharge on the day of surgery was reported as feasible in selected patients [190–192]. However, there was some disagreement around the safety of same-day discharge [193, 194]. The adoption and verification of a discharge checklist may be a useful tool for a safe discharge [57]. The minimum criteria for a discharge included vital parameters within the normal range, adequate pain control by nonopioid analgesics, adequate water intake, ability to tolerate liquid diet, and no evidence of sepsis or signs of postoperative complication. Finally, predischARGE education concerning signs and symptoms of possible postoperative complications and procedures for contacting the staff were also identified as a key component of a successful discharge [11, 12, 39, 55].

### Discussion

Literature review supports ERABS in reducing LOS without increasing complications and costs [8–10]. Producing a statement on ERABS standardizes the ERABS approach across bariatric surgery centers, benefiting both patients and hospitals [195]. The standardization of the approach may also allow the evaluation of ERABS's effective impact on the postoperative complications and total costs compared to the standard approach [196]. Notably, it became clear from this statement that the more items in the protocol that are adopted, the more efficient the ERABS approach will be [13].

Perioperative management involves a multidisciplinary team, where a full collaboration among anesthesiologists, surgeons, and staff members may lead to the best results [73]. The ERABS statement, drafted thanks to this collaboration, is, therefore, addressed not only to anesthesiologists or surgeons, but also to everyone involved in the perioperative care of obese patients in bariatric surgery centers.

**Acknowledgements** The authors acknowledge Medtronic as the sponsor of the meeting organized in order to realize the activities.

**Funding** Open access funding provided by Università degli Studi di Padova within the CRUI-CARE Agreement.

## Declarations

**Disclosures** Drs Giuseppe Marinari, Mirto Foletto, Carlo Nagliati, Giuseppe Navarra, Vincenzo Borrelli, Vincenzo Bruni, Giovanni Fantola, Roberto Moroni, Luigi Tritapepe, Roberta Monzani, Daniela Sanna, Michele Carron, Rita Cataldo have no conflicts of interest or financial ties to disclose.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Gloy VL, Briel M, Bhatt DL, Kashyap SR, Schauer PR, Mingrone G, Bucher HC, Nordmann AJ (2013) Bariatric surgery versus non-surgical treatment for obesity: a systematic review and meta-analysis of randomised controlled trials. *BMJ* 347:f5934
- Colquitt JL, Picot J, Loveman E, Clegg AJ (2009) Surgery for obesity. *Cochrane Database Syst Rev* 2:CD003641
- Smith TW Jr, Wang X, Singer MA, Godellas CV, Vaince FT (2020) Enhanced recovery after surgery: a clinical review of implementation across multiple surgical subspecialties. *Am J Surg* 219:530–534
- Thorell A, McCormick AD, Awad S, Reynolds N, Roulin D, Demartines N, Vignaud M, Alvarez A, Singh PM, Lobo DN (2016) Guidelines for perioperative care in bariatric surgery: Enhanced Recovery After Surgery (ERAS) society recommendations. *World J Surg* 40:2065–2083
- The Italian Ministry of Health (2018) Annual report on hospitalization activity. [http://www.salute.gov.it/imgs/C\\_17\\_pubblicazioni\\_2898\\_allegato.pdf](http://www.salute.gov.it/imgs/C_17_pubblicazioni_2898_allegato.pdf)
- OCEBM Levels of Evidence Working Group\*. The Oxford 2011 levels of evidence. <https://www.cebm.net/wp-content/uploads/2014/06/CEBM-Levels-of-Evidence-2.1.pdf>
- 7Grade Definitions. U.S. preventive services task force. October 2018. <https://uspreventiveservicestaskforce.org/uspstf/grade-definitions>
- Ahmed OS, Rogers AC, Bolger JC, Mastrosimone A, Robb WB (2018) Meta-analysis of enhanced recovery protocols in bariatric surgery. *J Gastrointest Surg* 22:964–972
- Singh PM, Panwar R, Borle A, Goudra B, Tripathi A, van Wageningen BA, Sinha A (2017) Efficiency and safety effects of applying ERAS protocols to bariatric surgery: a systematic review with meta-analysis and trial sequential analysis of evidence. *Obes Surg* 27:489–501
- Małczak P, Pisarska M, Piotr M, Wysocki M, Budzyński A, Pędziwiatr M (2017) Enhanced recovery after bariatric surgery: systematic review and meta-analysis. *Obes Surg* 27:226–235
- Lemanu DP, Singh PP, Berridge K, Burr M, Birch C, Babor R, McCormick AD, Arroll B, Hill AG (2013) Randomized clinical trial of enhanced recovery versus standard care after laparoscopic sleeve gastrectomy. *Br J Surg* 100:482–489
- Ruiz-Tovar J, Garcia A, Ferrigni C, Gonzalez J, Castellon C, Duran M (2019) Impact of implementation of an enhanced recovery after surgery (ERAS) program in laparoscopic Roux-en-Y gastric bypass: a prospective randomized clinical trial. *Surg Obes Relat Dis* 15:228–235
- Brethauer SA, Grieco A, Fraker T, Evans-Labok K, Smith A, McEvoy MD, Saber AA, Morton JM, Petrick A (2019) Employing Enhanced Recovery Goals in Bariatric Surgery (ENERGY): a national quality improvement project using the Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program. *Surg Obes Relat Dis* 15:1977–1989
- Geubbels N, Evren I, Acherman YIZ, Bruin SC, van de Laar AWJM, Hoen MB, de Brauw LM (2019) Randomized clinical trial of an enhanced recovery after surgery programme versus conventional care in laparoscopic Roux-en-Y gastric bypass surgery. *BJS Open* 3:274–281
- Simonelli V, Goergen M, Orlando GG, Arru L, Zolotas CA, Geeroms M, Poulain V, Azagra JS (2016) Fast-track in bariatric and metabolic surgery: feasibility and cost analysis through a matched-cohort study in a single centre. *Obes Surg* 26:1970–1977
- Elliott JA, Patel VM, Kirresh A, Ashrafian H, Le Roux CW, Olbers T, Athanasiou T, Zacharakis E (2013) Fast-track laparoscopic bariatric surgery: a systematic review. *Updates Surg* 65:85–94
- Deneuvy A, Slim K, Sodji M, Blanc P, Gallet D, Blanchet MC (2018) Implementation of enhanced recovery programs for bariatric surgery. Results from the Francophone large-scale database. *Surg Obes Relat Dis* 14:99–105
- Lam J, Suzuki T, Bernstein D, Zhao B, Maeda C, Pham T, Sandler BJ, Jacobsen GR, Cheverie JN, Horgan S (2019) An ERAS protocol for bariatric surgery: is it safe to discharge on post-operative day 1? *Surg Endosc* 33:580–586
- Mills E, Eyawo O, Lockhart I, Kelly S, Wu P, Ebbert JO (2011) Smoking cessation reduces postoperative complications: a systematic review and meta-analysis. *Am J Med* 124:144–154
- Carron M, Safaee Fakhr B, Iepariello G, Foletto M (2020) Perioperative care of the obese patient. *Br J Surg* 107:e39–e55
- Mechanick JI, Apovian C, Brethauer S, Garvey WT, Joffe AM, Kim J, Kushner RF, Lindquist R, Pessah-Pollack R, Seger J, Urman RD, Adams S, Cleek JB, Correa R, Figaro MK, Flanders K, Grams J, Hurley DL, Kothari S, Seger MV, Still CD (2020) Clinical practice guidelines for the perioperative nutrition, metabolic, and nonsurgical support of patients undergoing bariatric procedures—2019 update: cosponsored by American Association of Clinical Endocrinologists/American College of Endocrinology, The Obesity Society, American Society for Metabolic & Bariatric Surgery, Obesity Medicine Association, and American Society of Anesthesiologists. *Surg Obes Relat Dis* 16:175–247
- Edholm D, Kullberg J, Haenni A, Karlsson FA, Ahlström A, Hedberg J, Ahlström H, Sundbom M (2011) Preoperative 4-week low-calorie diet reduces liver volume and intrahepatic fat, and facilitates laparoscopic gastric bypass in morbidly obese. *Obes Surg* 21:345–350
- Di Lorenzo N, Antoniou SA, Batterham RL, Busetto L, Godoroja D, Iossa A, Carrano FM, Agresta F, Alarçon I, Azran C, Bouvy N, Balagué Ponz C, Buza M, Copescu C, De Luca M, Dicker D, Di Vincenzo A, Felsenreich DM, Francis NK, Fried M, Gonzalo Prats B, Goitein D, Halford JCG, Herlesova J, Kalogridaki M,

- Ket H, Morales-Conde S, Piatto G, Prager G, Pruijssers S, Pucci A, Rayman S, Romano E, Sanchez-Cordero S, Vilallonga R, Silecchia G (2020) Clinical practice guidelines of the European Association for Endoscopic Surgery (EAES) on bariatric surgery: update 2020 endorsed by IFSO-EC, EASO and ESPCOP. *Surg Endosc* 34:2332–2358
24. Anderin C, Gustafsson UO, Heijbel N, Thorell A (2015) Weight loss before bariatric surgery and postoperative complications: data from the Scandinavian Obesity Registry (SOREg). *Ann Surg* 261:909–913
25. Kaw R, Chung F, Pasupuleti V, Mehta J, Gay PC, Hernandez AV (2012) Meta-analysis of the association between obstructive sleep apnoea and postoperative outcome. *Br J Anaesth* 109:897–906
26. de Raaff CA, Coblijn UK, de Vries N, van Wagensveld BA (2016) Is fear for postoperative cardiopulmonary complications after bariatric surgery in patients with obstructive sleep apnea justified? A systematic review. *Am J Surg* 211:793–801
27. Goucham AB, Coblijn UK, Hart-Sweet HB, de Vries N, Lagarde SM, van Wagensveld BA (2016) Routine postoperative monitoring after bariatric surgery in morbidly obese patients with severe obstructive sleep Apnea: ICU admission is not necessary. *Obes Surg* 26:737–742
28. Corso R, Russotto V, Gregoretti C, Cattano D (2018) Perioperative management of obstructive sleep apnea: a systematic review. *Minerva Anestesiol* 84:81–93
29. Devaraj U, Rajagopala S, Kumar A, Ramachandran P, Devereaux PJ, D'Souza GA (2017) Undiagnosed obstructive sleep Apnea and postoperative outcomes: a prospective observational study. *Respiration* 94:18–25
30. Nepomnayshy D, Hesham W, Erickson B, MacDonald J, Iorio R, Brams D (2013) Sleep apnea: is routine preoperative screening necessary? *Obes Surg* 23:287–291
31. Romero-Corral A, Caples SM, Lopez-Jimenez F, Somers VK (2010) Interactions between obesity and obstructive sleep apnea: implications for treatment. *Chest* 137:711–719
32. Carron M, Zarantonello F, Tellaroli P, Ori C (2016) Perioperative noninvasive ventilation in obese patients: a qualitative review and meta-analysis. *Surg Obes Relat Dis* 12:681–691
33. Petrini F, Di Giacinto I, Cataldo R, Esposito C, Pavoni V, Donato P, Trollo A, Merli G, Sorbello M, Pelosi P, Obesity Task Force for the SIAARTI Airway Management Study Group (2016) Perioperative and periprocedural airway management and respiratory safety for the obese patient: 2016 SIAARTI Consensus. *Minerva Anestesiol* 82:1314–1335
34. Alvarez A, Goudra BG, Singh PM (2017) Enhanced recovery after bariatric surgery. *Curr Opin Anaesthesiol* 30:133–139
35. Grant MC, Gibbons MM, Ko CY, Wick EC, Cannesson M, Scott MJ, McEvoy MD, King AB, Wu CL (2019) Evidence review conducted for the agency for healthcare research and quality safety program for improving surgical care and recovery: focus on anesthesiology for bariatric surgery. *Anesth Analg* 129:51–60
36. Tsang E, Lambert E, Carey S (2018) Fasting leads to fasting: examining the relationships between perioperative fasting times and fasting for symptoms in patients undergoing elective abdominal surgery. *Asia Pac J Clin Nutr* 27:968–974
37. Lambert E, Carey S (2016) Practice guideline recommendations on perioperative fasting: a systematic review. *JPEN J Parenter Enteral Nutr* 40:1158–1165
38. Major P, Małczak P, Wysocki M, Torbic G, Gajewska N, Pędziwiatr M, Budzyński A (2018) Bariatric patients' nutritional status as a risk factor for postoperative complications, prolonged length of hospital stay and hospital readmission: a retrospective cohort study. *Int J Surg* 56:210–214
39. Trotta M, Ferrari C, D'Alessandro G, Sarra G, Piscitelli G, Marinari GM (2019) Enhanced recovery after bariatric surgery (ERABS) in a high-volume bariatric center. *Surg Obes Relat Dis* 15:1785–1792
40. Goretti G, Marinari GM, Vanni E, Ferrari C (2020) Value-based healthcare and enhanced recovery after surgery implementation in a high-volume bariatric center in Italy. *Obes Surg* 30:2519–2527
41. Prabhakaran S, Misra S, Magila M, Kumar SS, Kasthuri S, Palanivelu C, Raj PP (2020) Randomized controlled trial comparing the outcomes of enhanced recovery after surgery and standard recovery pathways in laparoscopic sleeve gastrectomy. *Obes Surg* 30:3273–3279
42. Pędziwiatr M, Mavriks J, Witowski J, Adamos A, Major P, Nowakowski M, Budzyński A (2018) Current status of enhanced recovery after surgery (ERAS) protocol in gastrointestinal surgery. *Med Oncol* 35:95
43. Gan TJ, Diemunsch P, Habib AS, Kovac A, Kranke P, Meyer TA, Watcha M, Chung F, Angus S, Apfel CC, Bergese SD, Candiotti KA, Chan MT, Davis PJ, Hooper VD, Lagoo-Deenadayalan S, Myles P, Nezat G, Philip BK, Tramèr MR, Society for Ambulatory Anesthesia (2014) Consensus guidelines for the management of postoperative nausea and vomiting. *Anesth Analg* 118:85–113
44. Benevides ML, Oliveira SS, de Aguilar-Nascimento JE (2013) The combination of haloperidol, dexamethasone, and ondansetron for prevention of postoperative nausea and vomiting in laparoscopic sleeve gastrectomy: a randomized double-blind trial. *Obes Surg* 23:1389–1396
45. Stein PD, Matta F (2013) Pulmonary embolism and deep venous thrombosis following bariatric surgery. *Obes Surg* 23:663–668
46. Gambhir S, Inaba CS, Alizadeh RF, Nahmias J, Hinojosa M, Smith BR, Nguyen NT, Daly S (2020) Venous thromboembolism risk for the contemporary bariatric surgeon. *Surg Endosc* 34:3521–3526
47. American Society for Metabolic and Bariatric Surgery Clinical Issues Committee (2013) ASMBS updated position statement on prophylactic measures to reduce the risk of venous thromboembolism in bariatric surgery patients. *Surg Obes Relat Dis* 9:493–497
48. Venclauskas L, Maleckas A, Arcelus JJ, ESA VTE Guidelines Task Force (2018) European guidelines on perioperative venous thromboembolism prophylaxis: surgery in the obese patient. *Eur J Anaesthesiol* 35:147–153
49. Bartlett MA, Mauck KF, Daniels PR (2015) Prevention of venous thromboembolism in patients undergoing bariatric surgery. *Vasc Health Risk Manag* 11:461–477
50. Ma P, Lloyd A, McGrath M, Moore R, Jackson A, Boone K, Higa K (2020) Reduction of opioid use after implementation of enhanced recovery after bariatric surgery (ERABS). *Surg Endosc* 34:2184–2190
51. Sinha A, Jayaraman L, Punhani D, Chowbey P (2017) Enhanced recovery after bariatric surgery in the severely obese, morbidly obese, super-morbidly obese and super-super morbidly obese using evidence-based clinical pathways: a comparative study. *Obes Surg* 27:560–568
52. Altieri MS, Yang J, Hajagos J, Spaniolas K, Park J, Gasparis AP, Bates AT, Docimo S, Talamini M, Shroyer AL, Pryor AD (2018) Evaluation of VTE prophylaxis and the impact of alternate regimens on post-operative bleeding and thrombotic complications following bariatric procedures. *Surg Endosc* 32:4805–4812
53. Raftopoulos I, Martindale C, Cronin A, Steinberg J (2008) The effect of extended post-discharge chemical thromboprophylaxis on venous thromboembolism rates after bariatric surgery: a prospective comparison trial. *Surg Endosc* 22:2384–2391

54. Dang JT, Szeto VG, Elnahas A, Ellsmere J, Okrainec A, Neville A, Malik S, Yorke E, Hong D, Biertho L, Jackson T, Karmali S (2020) Canadian Consensus Statement: enhanced recovery after surgery in bariatric surgery. *Surg Endosc* 34:1366–1375
55. Mannaerts GHH, Allatif REA, Al Hashmi FY, Bhosale A, Hammo AN, Isied SH, Qureshi WA, Al Hamad OS, Kayyal Y, Al Afari HST (2019) First successful large-scale introduction of an Enhanced Recovery after Bariatric Surgery (ERABS) Program in the Middle East: The Results and Lessons Learned of Tawam Hospital/Johns Hopkins, a Tertiary Governmental Center in the UAE. *Obes Surg* 29:2100–2109
56. Nagliati C, Troian M, Pennisi D, Balani A (2019) Enhanced recovery after bariatric surgery: 202 consecutive patients in an Italian Bariatric Center. *Obes Surg* 29:3133–3141
57. Gondal AB, Hsu CH, Serrot F, Rodriguez-Restrepo A, Hurbon AN, Galvani C, Ghaderi I (2019) Enhanced recovery in bariatric surgery: a study of short-term outcomes and compliance. *Obes Surg* 29:492–498
58. Awad S, Carter S, Purkayastha S, Hakky S, Moorthy K, Cousins J, Ahmed AR (2014) Enhanced recovery after bariatric surgery (ERABS): clinical outcomes from a tertiary referral bariatric centre. *Obes Surg* 24:753–758
59. Meunier H, Le Roux Y, Fiant AL, Marion Y, Bion AL, Gautier T, Contival N, Lubrano J, Fobe F, Zamparini M, Piquet MA, Savey V, Alves A, Menahem B (2019) Does the implementation of enhanced recovery after surgery (ERAS) guidelines improve outcomes of bariatric surgery? A propensity score analysis in 464 patients. *Obes Surg* 29:2843–2853
60. Major P, Wysocki M, Torbicz G, Gajewska N, Dudek A, Małczak P, Pędziwiatr M, Pisarska M, Radkowiak D, Budzyński A (2018) Risk factors for prolonged length of hospital stay and readmissions after laparoscopic sleeve gastrectomy and laparoscopic Roux-en-Y Gastric Bypass. *Obes Surg* 28:323–332
61. Proczko M, Kaska L, Twardowski P, Stepaniak P (2016) Implementing enhanced recovery after bariatric surgery protocol: a retrospective study. *J Anesth* 30:170–173
62. Fischer MI, Dias C, Stein A, Meinhardt NG, Heineck I (2014) Antibiotic prophylaxis in obese patients submitted to bariatric surgery. A systematic review. *Acta Cir Bras* 29:209–217
63. Bratzler DW, Dellinger EP, Olsen KM, Perl TM, Auwaerter PG, Bolon MK, Fish DN, Napolitano LM, Sawyer RG, Slain D, Steinberg JP, Weinstein RA, American Society of Health-System Pharmacists; Infectious Disease Society of America; Surgical Infection Society; Society for Healthcare Epidemiology of America (2013) Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Am J Health Syst Pharm* 70:195–283
64. Ferraz ÁAB, Santa-Cruz F, Edmiston CE Jr (2020) Antibiotic prophylaxis in bariatric surgical procedures: Is there an ideal antimicrobial agent? *Surg Infect (Larchmt)* 21:654–658
65. Chen X, Brathwaite CE, Barkan A, Hall K, Chu G, Cherasard P, Wang S, Nicolau DP, Islam S, Cunha BA (2017) Optimal cefazolin prophylactic dosing for bariatric surgery: no need for higher doses or intraoperative redosing. *Obes Surg* 27:626–629
66. Blum S, Cunha CB, Cunha BA (2019) Lack of pharmacokinetic basis of weight-based dosing and intra-operative re-dosing with cefazolin surgical prophylaxis in obese patients: implications for antibiotic stewardship. *Surg Infect (Larchmt)* 20:439–443
67. Hahl T, Peromaa-Haavisto P, Tarkkiainen P, Knutar O, Victorzon M (2016) Outcome of laparoscopic gastric bypass (LRYGB) with a program for enhanced recovery after surgery (ERAS). *Obes Surg* 26:505–511
68. Checketts MR, Alladi R, Ferguson K, Gemmell L, Handy JM, Klein AA, Love NJ, Misra U, Morris C, Nathanson MH, Rodney GE, Verma R, Pandit JJ, Association of Anaesthetists of Great Britain and Ireland (2016) Recommendations for standards of monitoring during anaesthesia and recovery 2015: Association of Anaesthetists of Great Britain and Ireland. *Anaesthesia* 71:85–93
69. Subramani Y, Riad W, Chung F, Wong J (2017) Optimal propofol induction dose in morbidly obese patients: a randomized controlled trial comparing the bispectral index and lean body weight scalar. *Can J Anaesth* 64:471–479
70. Gaszyński T, Wieczorek A (2016) A comparison of BIS recordings during propofol-based total intravenous anaesthesia and sevoflurane-based inhalational anaesthesia in obese patients. *Anaesthesiol Intensive Ther* 48:239–247
71. Blobner M, Hunter JM, Meistelman C, Hoeft A, Hollmann MW, Kirmeier E, Lewald H, Ulm K (2020) Use of a train-of-four ratio of 0.95 versus 0.9 for tracheal extubation: an exploratory analysis of POPULAR data. *Br J Anaesth* 124:63–72
72. Mason DS, Sapala JA, Wood MH, Sapala MA (1998) Influence of a forced air warming system on morbidly obese patients undergoing Roux-en-Y gastric bypass. *Obes Surg* 8:453–460
73. Carron M, Iepariello G, Safaee Fakhr B, Foletto M (2020) Perioperative care of the obese patient: a successful approach. *Br J Surg* 107:772–773
74. Leoni A, Arlati S, Ghisi D, Verweij M, Lugani D, Ghisi P, Cappelleri G, Cedrati V, El Tantswi Ali Alsheraei A, Pocar M, Ceriani V, Aldegheri G (2014) Difficult mask ventilation in obese patients: analysis of predictive factors. *Minerva Anestesiol* 80:149–157
75. Riad W, Vaez MN, Raveendran R, Tam AD, Quereshey FA, Chung F, Wong DT (2016) Neck circumference as a predictor of difficult intubation and difficult mask ventilation in morbidly obese patients: a prospective observational study. *Eur J Anaesthesiol* 33:244–249
76. Moon TS, Fox PE, Somasundaram A, Minhajuddin A, Gonzales MX, Pak TJ, Ogunnaike B (2019) The influence of morbid obesity on difficult intubation and difficult mask ventilation. *J Anesth* 33:96–102
77. Ravesloot MJ, van Maanen JP, Hilgevoord AA, van Wagenveld BA, de Vries N (2012) Obstructive sleep apnea is underrecognized and underdiagnosed in patients undergoing bariatric surgery. *Eur Arch Otorhinolaryngol* 269:1865–1871
78. Kapur VK, Auckley DH, Chowdhuri S, Kuhlmann DC, Mehra R, Ramar K, Harrod CG (2017) Clinical practice guideline for diagnostic testing for adult obstructive sleep Apnea: an American Academy of Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med* 13:479–504
79. de Raaff CA, Pierik AS, Coblijn UK, de Vries N, Bonjer HJ, van Wagenveld BA (2017) Value of routine polysomnography in bariatric surgery. *Surg Endosc* 31:245–248
80. Khan A, King WC, Patterson EJ, Laut J, Raum W, Courcoulas AP, Atwood C, Wolfe BM (2013) Assessment of obstructive sleep apnea in adults undergoing bariatric surgery in the longitudinal assessment of bariatric surgery-2 (LABS-2) study. *J Clin Sleep Med* 9:21–29
81. Lockhart EM, Willingham MD, Abdallah AB, Helsten DL, Bedair BA, Thomas J, Duntley S, Avidan MS (2013) Obstructive sleep apnea screening and postoperative mortality in a large surgical cohort. *Sleep Med* 14:407–415
82. Frerk C, Mitchell VS, McNarry AF, Mendonca C, Bhargava R, Patel A, O'Sullivan EP, Woodall NM, Ahmad I, Difficult Airway Society intubation guidelines working group (2015) Difficult Airway Society 2015 guidelines for management of unanticipated difficult intubation in adults. *Br J Anaesth* 115:827–848
83. Altermatt FR, Muñoz HR, Delfino AE, Cortés LI (2005) Pre-oxygenation in the obese patient: effects of position on tolerance to apnoea. *Br J Anaesth* 95:706–709
84. Guitton C, Ehrmann S, Volteau C, Colin G, Maamar A, Jean-Michel V, Mahe PJ, Landais M, Brule N, Bretonnière C,

- Zambon O, Vourc'h M (2019) Nasal high-flow preoxygenation for endotracheal intubation in the critically ill patient: a randomized clinical trial. *Intensive Care Med* 45:447–458
85. Vourc'h M, Huard D, Feuillet F, Baud G, Guichoux A, Surbled M, Tissot M, Chiffolleau A, Guitton C, Jaber S, Asehnoune K (2019) Preoxygenation in difficult airway management: high-flow oxygenation by nasal cannula versus face mask (the PRE-OPTIDAM study). Protocol for a single-centre randomised study. *BMJ Open* 9(4):e025909
  86. Cressey DM, Berthoud MC, Reilly CS (2001) Effectiveness of continuous positive airway pressure to enhance pre-oxygenation in morbidly obese women. *Anaesthesia* 56:680–684
  87. Coussa M, Proietti S, Schnyder P, Frascarolo P, Suter M, Spahn DR, Magnusson L (2004) Prevention of atelectasis formation during the induction of general anesthesia in morbidly obese patients. *Anesth Analg* 98:1491–1495
  88. Gander S, Frascarolo P, Suter M, Spahn DR, Magnusson L (2005) Positive end-expiratory pressure during induction of general anesthesia increases duration of nonhypoxic apnea in morbidly obese patients. *Anesth Analg* 100:580–584
  89. Edmark L, Östberg E, Scheer H, Wallquist W, Hedenstierna G, Zetterström H (2016) Preserved oxygenation in obese patients receiving protective ventilation during laparoscopic surgery: a randomized controlled study. *Acta Anaesthesiol Scand* 60:26–35
  90. Georgescu M, Tanoubi I, Fortier LP, Donati F, Drolet P (2012) Efficacy of preoxygenation with non-invasive low positive pressure ventilation in obese patients: crossover physiological study. *Ann Fr Anesth Reanim* 31:e161–e165
  91. Harbut P, Gozdzik W, Stjernfält E, Marsk R, Hesselvik JF (2014) Continuous positive airway pressure/pressure support pre-oxygenation of morbidly obese patients. *Acta Anaesthesiol Scand* 58:675–680
  92. Hoshijima H, Denawa Y, Tominaga A, Nakamura C, Shiga T, Nagasaka H (2018) Videolaryngoscope versus Macintosh laryngoscope for tracheal intubation in adults with obesity: a systematic review and meta-analysis. *J Clin Anesth* 44:69–75
  93. Carron M, Iepariello G, Linassi F (2021) Videolaryngoscopy versus direct laryngoscopy for tracheal intubation in obese adults: a meta-analysis. *J Clin Anesth* 71:110216
  94. Sinha A, Jayaraman L, Punhani D (2013) ProSeal™ LMA increases safe apnea period in morbidly obese patients undergoing surgery under general anesthesia. *Obes Surg* 23:580–584
  95. Natalini G, Franceschetti ME, Pantelidi MT, Rosano A, Lanza G, Bernardini A (2003) Comparison of the standard laryngeal mask airway and the ProSeal laryngeal mask airway in obese patients. *Br J Anaesth* 90:323–326
  96. Cook TM, Woodall N, Frerk C, Fourth National Audit Project (2011) Major complications of airway management in the UK: results of the Fourth National Audit Project of the royal college of anaesthetists and the difficult airway society. Part 1: anaesthesia. *Br J Anaesth* 106:617–631
  97. Cook TM, Woodall N, Harper J, Benger J, Fourth National Audit Project (2011) Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the Difficult Airway Society. Part 2: intensive care and emergency departments. *Br J Anaesth* 106:632–642
  98. Nicholson A, Cook TM, Smith AF, Lewis SR, Reed SS (2013) Supraglottic airway devices versus tracheal intubation for airway management during general anaesthesia in obese patients. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD010105.pub2>
  99. Ahmad I, El-Boghdady K, Bhagrath R, Hodzovic I, McNarry AF, Mir F, O'Sullivan EP, Patel A, Stacey M, Vaughan D (2020) Difficult Airway Society guidelines for awake tracheal intubation (ATI) in adults. *Anaesthesia* 75:509–528
  100. Abdelmalak BB, Bernstein E, Egan C, Abdallah R, You J, Sessler DI, Doyle DJ (2011) GlideScope® vs flexible fibreoptic scope for elective intubation in obese patients. *Anaesthesia* 66:550–555
  101. Rosenstock CV, Thøgersen B, Afshari A, Christensen AL, Eriksen C, Gätke MR (2012) Awake fiberoptic or awake video laryngoscopic tracheal intubation in patients with anticipated difficult airway management: a randomized clinical trial. *Anesthesiology* 116:1210–1216
  102. Abdellatif AA, Ali MA (2014) GlideScope videolaryngoscope versus flexible fiberoptic bronchoscope for awake intubation of morbidly obese patient with predicted difficult intubation. *Middle East J Anaesthesiol* 22:385–392
  103. Juvin P, Vadam C, Malek L, Dupont H, Marmuse JP, Desmots JM (2000) Postoperative recovery after desflurane, propofol, or isoflurane anesthesia among morbidly obese patients: a prospective, randomized study. *Anesth Analg* 91:714–719
  104. La Colla L, Albertin A, La Colla G, Mangano A (2007) Faster wash-out and recovery for desflurane vs sevoflurane in morbidly obese patients when no premedication is used. *Br J Anaesth* 99:353–358
  105. McKay RE, Malhotra A, Cakmakaya OS, Hall KT, McKay WR, Apfel CC (2010) Effect of increased body mass index and anaesthetic duration on recovery of protective airway reflexes after sevoflurane vs desflurane. *Br J Anaesth* 104:175–182
  106. Liu FL, Chong YG, Chen SY, Su YH, Huang SY, Lo PH, Lee YY, Tam KW (2015) Postoperative recovery after anesthesia in morbidly obese patients: a systematic review and meta-analysis of randomized controlled trials. *Can J Anaesth* 62:907–917
  107. Sneyd JR, Carr A, Byrom WD, Bilski AJ (1998) A meta-analysis of nausea and vomiting following maintenance of anaesthesia with propofol or inhalational agents. *Eur J Anaesthesiol* 15:433–445
  108. Carron M, Tessari I, Linassi F, Navalesi P (2021) Desflurane versus propofol for general anesthesia maintenance in obese patients: a pilot meta-analysis. *J Clin Anesth* 68:110103
  109. De Baerdemaeker L, Margaron M (2016) Best anaesthetic drug strategy for morbidly obese patients. *Curr Opin Anaesthesiol* 29:119–128
  110. Frauenknecht J, Kirkham KR, Jacot-Guillarmod A, Albrecht E (2019) Analgesic impact of intra-operative opioids vs. opioid-free anaesthesia: a systematic review and meta-analysis. *Anaesthesia* 74:651–662
  111. Ehsan Z, Mahmoud M, Shott SR, Amin RS, Ishman SL (2016) The effects of anesthesia and opioids on the upper airway: a systematic review. *Laryngoscope* 126:270–284
  112. Ziemann-Gimmel P, Goldfarb AA, Koppman J, Marema RT (2014) Opioid-free total intravenous anaesthesia reduces post-operative nausea and vomiting in bariatric surgery beyond triple prophylaxis. *Br J Anaesth* 112:906–911
  113. Mulier JP, Dillemans B (2019) Anaesthetic factors affecting outcome after bariatric surgery, a retrospective levelled regression analysis. *Obes Surg* 29:1841–1850
  114. Komatsu R, Turan AM, Orhan-Sungur M, McGuire J, Radke OC, Apfel CC (2007) Remifentanyl for general anaesthesia: a systematic review. *Anaesthesia* 62:1266–1280
  115. Sudré EC, de Batista PR, Castiglia YM (2015) Longer immediate recovery time after anesthesia increases risk of respiratory complications after laparotomy for bariatric surgery: a randomized clinical trial and a cohort study. *Obes Surg* 25:2205–2212
  116. Choi YK, Brolin RE, Wagner BK, Chou S, Etesham S, Pollak P (2000) Efficacy and safety of patient-controlled analgesia for morbidly obese patients following gastric bypass surgery. *Obes Surg* 10:154–159
  117. Andersen LP, Werner MU, Rosenberg J, Gögenur I (2014) Analgesic treatment in laparoscopic gastric bypass surgery: a systematic review of randomized trials. *Obes Surg* 24:462–470



118. Ng JJ, Leong WQ, Tan CS, Poon KH, Lomanto D, So JBY, Shabbir A (2017) Multimodal analgesic protocol reduces opioid-related adverse events and improves patient outcomes in laparoscopic sleeve gastrectomy. *Obes Surg* 27:3075–3081
119. Sollazzi L, Modesti C, Vitale F, Sacco T, Ciocchetti P, Idra AS, Tacchino RM, Perilli V (2009) Preinductive use of clonidine and ketamine improves recovery and reduces postoperative pain after bariatric surgery. *Surg Obes Relat Dis* 5:67–71
120. Kasputytė G, Karbonskienė A, Macas A, Maleckas A (2020) Role of ketamine in multimodal analgesia protocol for bariatric surgery. *Medicine (Kaunas)* 56:96
121. Naja ZM, Khatib R, Ziade FM, Moussa G, Naja ZZ, Naja AS, Kanawati S (2014) Effect of clonidine versus dexmedetomidine on pain control after laparoscopic gastric sleeve: a prospective, randomized, double-blinded study. *Saudi J Anaesth* 8(Suppl 1):S57–62
122. Bakhamees HS, El-Halafawy YM, El-Kerdawy HM, Gouda NM, Altemyatt S (2007) Effects of dexmedetomidine in morbidly obese patients undergoing laparoscopic gastric bypass. *Middle East J Anaesthesiol* 19:537–551
123. Zarif P, Abdelaal Ahmed Mahmoud A, Abdelhaq MM, Mikhail HM, Farag A (2016) Dexmedetomidine versus magnesium sulfate as adjunct during anesthesia for laparoscopic colectomy. *Anesthesiol Res Pract* 2016:7172920
124. El Mourad MB, Arafa SK (2019) Effect of intravenous versus intraperitoneal magnesium sulfate on hemodynamic parameters and postoperative analgesia during laparoscopic sleeve gastrectomy—a prospective randomized study. *J Anaesthesiol Clin Pharmacol* 35:242–247
125. Kizilcik N, Koner O (2018) Magnesium sulfate reduced opioid consumption in obese patients undergoing sleeve gastrectomy: a prospective, randomized clinical trial. *Obes Surg* 28:2783–2788
126. De Oliveira GS, Duncan K, Fitzgerald P, Nader A, Gould RW, McCarthy RJ (2014) Systemic lidocaine to improve quality of recovery after laparoscopic bariatric surgery: a randomized double-blinded placebo-controlled trial. *Obes Surg* 24:212–218
127. Salama AK, Abdallah NM (2016) Multimodal analgesia with pregabalin and dexmedetomidine in morbidly obese patients undergoing laparoscopic sleeve gastrectomy: a prospective randomized double blind placebo controlled study. *Egypt J Anaesth* 32:293–298
128. Alimian M, Imani F, Faiz SH, Pournajafian A, Navadegi SF, Safari S (2012) Effect of oral pregabalin premedication on post-operative pain in laparoscopic gastric bypass surgery. *Anesth Pain Med* 2:12–16
129. Rupniewska-Ladyko A, Malec-Milewska M, Kraszewska E, Pirozynski M (2018) Gabapentin before laparoscopic sleeve gastrectomy reduces postoperative oxycodone consumption in obese patients: a randomized double-blind placebo-controlled trial. *Minerva Anesthesiol* 84:565–571
130. Alkhamesi NA, Kane JM, Guske PJ, Wallace JW, Rantis PC (2008) Intraperitoneal aerosolization of bupivacaine is a safe and effective method in controlling postoperative pain in laparoscopic Roux-en-Y gastric bypass. *J Pain Res* 1:9–13
131. Omar I, Abuelsel A (2019) Efficacy of intraperitoneal instillation of bupivacaine after bariatric surgery: randomized controlled trial. *Obes Surg* 29:1735–1741
132. Safari S, Rokhtabnak F, Djalali Motlagh S, Ghanbari Garkani M, Pournajafian A (2020) Effect of intraperitoneal bupivacaine on postoperative pain in laparoscopic bariatric surgeries. *Surg Obes Relat Dis* 16:299–305
133. Ruiz-Tovar J, Gonzalez J, Garcia A, Cruz C, Rivas S, Jimenez M, Ferrigni C, Duran M (2016) Intraperitoneal ropivacaine irrigation in patients undergoing bariatric surgery: a prospective randomized clinical trial. *Obes Surg* 26:2616–2621
134. Albrecht E, Kirkham KR, Endersby RV, Chan VW, Jackson T, Okrainec A, Penner T, Jin R, Brull R (2013) Ultrasound-guided transversus abdominis plane (TAP) block for laparoscopic gastric-bypass surgery: a prospective randomized controlled double-blinded trial. *Obes Surg* 23:1309–1314
135. Ruiz-Tovar J, Gonzalez G, Sarmiento A, Carbajo MA, Ortiz-de-Solorzano J, Castro MJ, Jimenez JM, Zubiaga L (2020) Analgesic effect of postoperative laparoscopic-guided transversus abdominis plane (TAP) block, associated with preoperative port-site infiltration, within an enhanced recovery after surgery protocol in one-anastomosis gastric bypass: a randomized clinical trial. *Surg Endosc* 34:5455–5460
136. Mittal T, Dey A, Siddhartha R, Nali A, Sharma B, Malik V (2018) Efficacy of ultrasound-guided transversus abdominis plane (TAP) block for postoperative analgesia in laparoscopic gastric sleeve resection: a randomized single blinded case control study. *Surg Endosc* 32:4985–4989
137. Carron M, De Cassai A, Linassi F, Navalesi P (2020) Multimodal analgesia in bariatric surgery: not just an intravenous approach. *Surg Obes Relat Dis* 16:2133–2135
138. Hamid HKS, Ahmed AY, Saber AA, Emile SH, Ibrahim M, Ruiz-Tovar J (2020) Transversus abdominis plane block using a short-acting local anesthetic reduces pain and opioid consumption after laparoscopic bariatric surgery: a meta-analysis. *Surg Obes Relat Dis* 16:1349–1357
139. Ruiz-Tovar J, Muñoz JL, Gonzalez J, Zubiaga L, García A, Jimenez M, Ferrigni C, Durán M (2017) Postoperative pain after laparoscopic sleeve gastrectomy: comparison of three analgesic schemes (isolated intravenous analgesia, epidural analgesia associated with intravenous analgesia and port-sites infiltration with bupivacaine associated with intravenous analgesia). *Surg Endosc* 31:231–236
140. Fuchs-Buder T, Schmartz D, Baumann C, Hilt L, Nomine-Criqui C, Meistelman C, Brunaud L (2019) Deep neuromuscular blockade improves surgical conditions during gastric bypass surgery for morbid obesity: a randomised controlled trial. *Eur J Anaesthesiol* 36:486–493
141. Torensmas B, Martini CH, Boon M, Olofsen E, In't Veld B, Liem RS, Knook MT, Swank DJ, Dahan A (2016) Deep neuromuscular block improves surgical conditions during bariatric surgery and reduces postoperative pain: a randomized double blind controlled trial. *PLoS One* 11:e0167907
142. Gaszynski T, Szweczyk T, Gaszynski W (2012) Randomized comparison of sugammadex and neostigmine for reversal of rocuronium-induced muscle relaxation in morbidly obese undergoing general anaesthesia. *Br J Anaesth* 108:236–239
143. Carron M, Veronese S, Foletto M, Ori C (2013) Sugammadex allows fast-track bariatric surgery. *Obes Surg* 23:1558–1563
144. Castro DS Jr, Leão P, Borges S, Gomes L, Pacheco M, Figueiredo P (2014) Sugammadex reduces postoperative pain after laparoscopic bariatric surgery: a randomized trial. *Surg Laparosc Endosc Percutan Tech* 24:420–423
145. Writing Committee for the PROBESE Collaborative Group of the PROtective VEntilation Network (PROVENet) for the Clinical Trial Network of the European Society of Anaesthesiology, Bluth T, Serpa Neto A, Schultz MJ, Pelosi P, Gama de Abreu M; PROBESE Collaborative Group, Bluth T, Bobek I, Canet JC, Cinnella G, de Baerdemaeker L, Gama de Abreu M, Gregoretti C, Hedenstierna G, Hemmes SNT, Hiesmayr M, Hollmann MW, Jaber S, Laffey J, Licker MJ, Markstaller K, Matot I, Mills GH, Mulier JP, Pelosi P, Putensen C, Rossaint R, Schmitt J, Schultz MJ, Senturk M, Serpa Neto A, Severgnini P, Sprung J, Vidal Melo MF, Wrigge H (2019) Effect of intraoperative high positive end-expiratory pressure (PEEP) with recruitment maneuvers vs low PEEP on postoperative pulmonary complications in obese patients: a randomized clinical trial. *JAMA* 321:2292–2305



146. Ball L, Hemmes SNT, Serpa Neto A, Bluth T, Canet J, Hiesmayr M, Hollmann MW, Mills GH, Vidal Melo MF, Putensen C, Schmid W, Severgnini P, Wrigge H, Gama de Abreu M, Schultz MJ, Pelosi P, LAS VEGAS investigators; PROVE Network; Clinical Trial Network of the European Society of Anaesthesiology (2018) Intraoperative ventilation settings and their associations with postoperative pulmonary complications in obese patients. *Br J Anaesth* 121:899–908
147. O'Gara B, Talmor D (2018) Perioperative lung protective ventilation. *BMJ* 362(8222):k3030
148. Nguyen NT, Wolfe BM (2005) The physiologic effects of pneumoperitoneum in the morbidly obese. *Ann Surg* 241:219–226
149. Atkinson TM, Giraud GD, Togioka BM, Jones DB, Cigarroa JE (2017) Cardiovascular and ventilatory consequences of laparoscopic surgery. *Circulation* 135:700–710
150. Costa Souza GM, Santos GM, Zimpel SA, Melnik T (2020) Intraoperative ventilation strategies for obese patients undergoing bariatric surgery: systematic review and meta-analysis. *BMC Anesthesiol* 20:36
151. Schol PB, Terink IM, Lancé MD, Scheepers HC (2016) Liberal or restrictive fluid management during elective surgery: a systematic review and meta-analysis. *J Clin Anesth* 35:26–39
152. Pösö T, Kesek D, Aroch R, Winsö O (2013) Morbid obesity and optimization of preoperative fluid therapy. *Obes Surg* 23:1799–1805
153. Schuster R, Alami RS, Curet MJ, Paulraj N, Morton JM, Brodsky JB, Brock-Utne JG, Lemmens HJ (2006) Intra-operative fluid volume influences postoperative nausea and vomiting after laparoscopic gastric bypass surgery. *Obes Surg* 16:848–851
154. Brienza N, Biancofiore G, Cavaliere F, Corcione A, De Gasperi A, De Rosa RC, Fumagalli R, Giglio MT, Locatelli A, Lorini FL, Romagnoli S, Scolletta S, Tritapepe L (2019) Clinical guidelines for perioperative hemodynamic management of non-cardiac surgical adult patients. *Minerva Anestesiol* 85:1315–1333
155. Demirel İ, Bolat E, Altun AY, Özdemir M, Beştaş A (2018) Efficacy of goal-directed fluid therapy via pleth variability index during laparoscopic Roux-en-Y gastric bypass surgery in morbidly obese patients. *Obes Surg* 28:358–363
156. Jain AK, Dutta A (2010) Stroke volume variation as a guide to fluid administration in morbidly obese patients undergoing laparoscopic bariatric surgery. *Obes Surg* 20:709–715
157. Muñoz JL, Gabaldón T, Miranda E, Berrio DL, Ruiz-Tovar J, Ronda JM, Esteve N, Arroyo A, Pérez A (2016) Goal-directed fluid therapy on laparoscopic sleeve gastrectomy in morbidly obese patients. *Obes Surg* 26:2648–2653
158. Practice guidelines for preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration: application to healthy patients undergoing elective procedures: an updated report by the American society of Anesthesiologists task force on preoperative fasting and the use of pharmacologic agents to reduce the risk of pulmonary aspiration (2017) *Anesthesiology* 126:376–393
159. Salmasi V, Maheshwari K, Yang D, Mascha EJ, Singh A, Sessler DI, Kurz A (2017) Relationship between intraoperative hypotension, defined by either reduction from baseline or absolute thresholds, and acute kidney and myocardial injury after non-cardiac surgery: a retrospective cohort analysis. *Anesthesiology* 126:47–65
160. Difficult Airway Society Extubation Guidelines Group, Popat M, Mitchell V, Dravid R, Patel A, Swamipillai C, Higgs A (2012) Difficult Airway Society Guidelines for the management of tracheal extubation. *Anaesthesia* 67:318–340
161. Members of the Working Party, Nightingale CE, Margaron MP, Shearer E, Redman JW, Lucas DN, Cousins JM, Fox WT, Kennedy NJ, Venn PJ, Skues M, Gabbott D, Misra U, Pandit JJ, Popat MT, Griffiths R; Association of Anaesthetists of Great Britain; Ireland Society for Obesity and Bariatric Anaesthesia (2015) Peri-operative management of the obese surgical patient 2015: Association of Anaesthetists of Great Britain and Ireland Society for Obesity and Bariatric Anaesthesia. *Anaesthesia* 70:859–876
162. Rochweg B, Einav S, Chaudhuri D, Mancebo J, Mauri T, Helviz Y, Goligher EC, Jaber S, Ricard JD, Rittayamai N, Roca O, Antonelli M, Maggiore SM, Demoule A, Hodgson CL, Mercat A, Wilcox ME, Granton D, Wang D, Azoulay E, Ouannes-Besbes L, Cinnella G, Rauseo M, Carvalho C, Dessap-Mekontso A, Fraser J, Frat JP, Gomersall C, Grasselli G, Hernandez G, Jog S, Pesenti A, Riviello ED, Slutsky AS, Stapleton RD, Talmor D, Thille AW, Brochard L, Burns KEA (2020) The role for high flow nasal cannula as a respiratory support strategy in adults: a clinical practice guideline. *Intensive Care Med* 46:2226–2237
163. Zarembo S, Shin CH, Hutter MM, Malviya SA, Grabitz SD, MacDonald T, Diaz-Gil D, Ramachandran SK, Hess D, Malhotra A, Eikermann M (2016) Continuous positive airway pressure mitigates opioid-induced worsening of sleep-disordered breathing early after bariatric surgery. *Anesthesiology* 125:92–104
164. Nelson R, Edwards S, Tse B (2007) Prophylactic nasogastric decompression after abdominal surgery. *Cochrane Database Syst Rev* 2007:CD004929
165. Petrowsky H, Demartines N, Rousson V, Clavien PA (2004) Evidence-based value of prophylactic drainage in gastrointestinal surgery: a systematic review and meta-analyses. *Ann Surg* 240:1074–1084
166. Liscia G, Scaringi S, Facchiano E, Quartararo G, Lucchese M (2014) The role of drainage after Roux-en-Y gastric bypass for morbid obesity: a systematic review. *Surg Obes Relat Dis* 10:171–176
167. Kavuturu S, Rogers AM, Haluck RS (2012) Routine drain placement in Roux-en-Y gastric bypass: an expanded retrospective comparative study of 755 patients and review of the literature. *Obes Surg* 22:177–181
168. Hoehn RS, Seitz AP, Singer KE, Thompson JR, Watkins BM (2019) Enhanced recovery protocol for laparoscopic sleeve gastrectomy: Are narcotics necessary? *J Gastrointest Surg* 23:1541–1546
169. Ronellenfitsch U, Schwarzbach M, Kring A, Kienle P, Post S, Hasenberg T (2012) The effect of clinical pathways for bariatric surgery on perioperative quality of care. *Obes Surg* 22:732–739
170. Saint S, Lipsky BA (1999) Preventing catheter-related bacteriuria: should we? Can we? How? *Arch Intern Med* 159:800–808
171. Kleven RM, Edwards JR, Richards CL Jr, Horan TC, Gaynes RP, Pollock DA, Cardo DM (2007) Estimating health care-associated infections and deaths in U.S. hospitals, 2002. *Public Health Rep* 122:160–166
172. Taylor J, Canner J, Cronauer C, Prior D, Coker A, Nguyen H, Magnuson T, Adrales G, Schweitzer M (2020) Implementation of an enhanced recovery program for bariatric surgery. *Surg Endosc* 34:2675–2681
173. Małczak P, Wysocki M, Twardowska H, Dudek A, Tabiś J, Major P, Pisarska M, Pędziwiatr M (2020) Impact of adherence to the ERAS® protocol on short-term outcomes after bariatric surgery. *Obes Surg* 30:1498–1505
174. Ljungqvist O, Scott M, Fearon KC (2017) Enhanced recovery after surgery: a review. *JAMA Surg* 152:292–298
175. Major P, Stefura T, Małczak P, Wysocki M, Witowski J, Kulawik J, Wierdak M, Pisarska M, Pędziwiatr M, Budzyński A (2018) Postoperative care and functional recovery after laparoscopic sleeve gastrectomy vs. laparoscopic Roux-en-Y gastric bypass among patients under ERAS protocol. *Obes Surg* 28:1031–1039
176. Balla A, Batista Rodríguez G, Corradetti S (2017) Outcomes after bariatric surgery according to large databases: a systematic review. *Langenbecks Arch Surg* 402:885–899

177. Ardila-Gatas J, Sharma G, Lloyd SJ, Khorgami Z, Tu C, Schauer PR, Brethauer SA, Aminian A (2019) A nationwide safety analysis of discharge on the first postoperative day after bariatric surgery in selected patients. *Obes Surg* 29:15–22
178. Chiappetta S, Jamadar P, Stier C, Bottino V, Weiner RA, Runkel N (2020) The role of C-reactive protein after surgery for obesity and metabolic disorders. *Surg Obes Relat Dis* 16:99–108
179. Muñoz JL, Ruiz-Tovar J, Miranda E, Berrio DL, Moya P, Gutiérrez M, Flores R, Picó C, Pérez A (2016) C-Reactive protein and procalcitonin as early markers of septic complications after laparoscopic sleeve gastrectomy in morbidly obese patients within an enhanced recovery after surgery program. *J Am Coll Surg* 222:831–837
180. Bona D, Micheletto G, Bonitta G, Panizzo V, Cavalli M, Rausa E, Cirri S, Aiolfi A (2019) Does C-reactive protein have a predictive role in the early diagnosis of postoperative complications after bariatric surgery? Systematic review and Bayesian meta-analysis. *Obes Surg* 29:3448–3456
181. Lee Y, McKechnie T, Doumouras AG, Handler C, Eskicioglu C, Gmora S, Anvari M, Hong D (2019) Diagnostic value of C-reactive protein levels in postoperative infectious complications after bariatric surgery: a systematic review and meta-analysis. *Obes Surg* 29:2022–2029
182. Williams M, McMeekin S, Wilson J, Miller G, Langlands F, Wong W, Peter M, Giles M (2017) Predictive value of C-reactive protein for complications post-laparoscopic Roux-En-Y Gastric bypass-author reply. *Obes Surg* 27:2462–2463
183. Kassir R, Blanc P, Bruna Tibalbo LM, Breton C, Lointier P (2015) C-Reactive protein and procalcitonin for the early detection of postoperative complications after sleeve gastrectomy: preliminary study in 97 patients. *Surg Endosc* 29:1439–1444
184. Williams MR, McMeekin S, Wilson RJ, Miller GV, Langlands FE, Wong W, Peter M, Giles MS (2017) Predictive value of C-reactive protein for complications post-laparoscopic Roux-En-Y Gastric bypass. *Obes Surg* 27:709–715
185. Williams M, Langlands F, Giles M (2020) Does C-reactive protein day 1 post-surgery have a predictive role for post-operative complications: a single-centre perspective following published meta-analysis. *Obes Surg* 30:347–348
186. Rebibo L, Cosse C, Robert B, Chivot C, Yzet T, Dhahri A, Regimbeau JM (2017) Eliminating routine upper gastrointestinal contrast studies after sleeve gastrectomy decreases length of stay and hospitalization costs. *Surg Obes Relat Dis* 13:553–559
187. Rahman U, Docimo S, Pryor AD, Bates A, Obeid NR, Spaniolas K (2018) Routine contrast imaging after bariatric surgery and the effect on hospital length of stay. *Surg Obes Relat Dis* 14:517–520
188. Diaz Vico T, Elli EF (2019) Utility of immediate postoperative upper gastrointestinal contrast study in bariatric surgery. *Obes Surg* 29:1130–1133
189. Brockmeyer JR, Simon TE, Jacob RK, Husain F, Choi Y (2012) Upper gastrointestinal swallow study following bariatric surgery: institutional review and review of the literature. *Obes Surg* 22:1039–1043
190. Quartararo G, Facchiano E, Scaringi S, Liscia G, Lucchese M (2014) Upper gastrointestinal series after Roux-en-Y gastric bypass for morbid obesity: effectiveness in leakage detection. a systematic review of the literature. *Obes Surg* 24:1096–1101
191. Leepalao MC, Arredondo D, Speights F, Duncan TD (2020) Same-day discharge on laparoscopic Roux-en-Y gastric bypass patients: an outcomes review. *Surg Endosc* 34:3614–3617
192. Aftab H, Fagerland MW, Gondal G, Ghanima W, Olsen MK, Nordby T (2019) Gastric sleeve resection as day-case surgery: what affects the discharge time? *Surg Obes Relat Dis* 15:2018–2024
193. Inaba CS, Koh CY, Sujatha-Bhaskar S, Pejcinovska M, Nguyen NT (2018) How safe is same-day discharge after laparoscopic sleeve gastrectomy? *Surg Obes Relat Dis* 14:1448–1453
194. Inaba CS, Koh CY, Sujatha-Bhaskar S, Zhang L, Nguyen NT (2018) Same-day discharge after laparoscopic Roux-en-Y gastric bypass: an analysis of the metabolic and bariatric surgery accreditation and quality improvement program database. *J Am Coll Surg* 226:868–873
195. Leeman M, van Mil SR, Biter LU, Apers JA, Verhoef K, Dunkelgrun M (2021) Reducing complication rates and hospital readmissions while revising the enhanced recovery after bariatric surgery (ERABS) protocol. *Surg Endosc* 35:612–619
196. Patil S, Cornett EM, Jesunathadas J, Belani K, Fox CJ, Kaye AD, Lambert LA, Urman RD (2019) Implementing enhanced recovery pathways to improve surgical outcomes. *J Anaesthesiol Clin Pharmacol* 35(Suppl 1):S24–S28

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.