

21-22 Dicembre 2020
XXVIII Congresso
Nazionale
SICOB ONLINE

Un palloncino li solleverà

RD. BERTA

U.O. Chirurgia Bariatrica e Metabolica AOU Pisa

Gastric Balloons Offer an Option for Patients who:

- Have failed attempts of medical weight loss
- need rapid minimal weight loss (for knee replacement, to enable fertility in patients with PCOS)
- Are out of indication for surgery (BMI< 35)

- Refuse Bariatric Surgery
- ARE SUPEROBESE WAITING FOR BARIATRIC SURGERY



opic Gastric B

NO INDICATION BEFORE BARIATRIC SURGERY

deposition in the pharynge

onary I

uce functional residual capacity

OBES SURG DOI 10.1007/s11695-016-2383-x

ORIGINAL CONTRIBUTIONS

Impact of Intragastric Balloon Before
Bypass on Patients with Super Obesity:
Multicenter Study

B. Coffin ¹² © &V. Maunoury ^{3,4} &F. Pattou ^{3,4} &X. Hébuterne ^{5,6} &S. Schneider ^{5,6} &M. Coupaye ^{1,2} &S. Ledoux ^{1,2} &F. Iglicki ^{1,2} &F. Mion ^{7,8} &M. Robert ^{7,8} &E. Disse ^{7,8} &J. Escourrou ^{9,10} &G. Tuyeras ^{9,10} &Y. Le Roux ¹¹ &C. Arvieux ^{12,13} &P. Pouderoux ^{14,15} &N. Huten ^{16,17} &T. Alfaiate ¹⁸ &D. Haiace ¹⁸ &S. Msika ^{1,2}

IGB insertion the perioperative

«3 severe compli

ed weight loss but did not improve st postoperative weight loss.

at

ccurred during IGB removal»

New Swallowable Intragastric Ballon: Elipse

No Need for Anesthesia /Sedation and Endoscopy



Aprepitant / Netupitant Granisetron / Palonsetron

OUTPATIENT SETTING

35.000 implants

Elipse Programme

Health Care Team - Smart Scale - App







After approximately 16 weeks, a time-activated release valve will open, allowing the balloon to empty and pass naturally through the gastrointestinal tract

EXCLUSION CRITERIA

- Difficulty swallowing
- Large hiatal hernia
- Active ulcer in the stomach or duodenum
- Previous surgery of the stomach
- Inflammatory bowel disease
- Variceal disease
- Alcoholic disease or drug abuse
- Psychiatric disease
- Pregnancy
- Use of anti-coagulants or anti-inflammatory drugs
- Cardiovascular, pulmonary or cerebrovascular disease
- Condition that predispose to bowel obstruction




Intragastric Balloons - Results from Pivotal RCTs




F Bazerbachi et al. Am J Gastroenterol 2019


Elipse COMPLICATIONS

Adverse Events	Elipse %	Others %
Premature deflation	0.50	0.90
Intolerance leading to early balloon removal	0.30	2.2
Small bowel obstruction	0.15	0.8
Hyperinflation	0.07	0.5
Gastric Perforation	0.03	0.2
Perforation during endoscopy		0.40
Incomplete device empting	0.03	
Pancreatitis	0.01	
Death	0	0.03









CONCLUSIONS

- Elipse can be used before bariatric surgery when attempts at medical weight loss management have failed
- Despite its semplicity and low invasiveness a multidisciplinary aproach is mandatory for patients' selection and prompt detection of complications that should be managed in an appropriate setting only by an expert bariatric team
- it is not covered by the national health system